

Macroscopic and Microscopic Description of Black Dipoles

Roberto Emparan^{a1} and Edward Teo^b

^a*Theory Division, CERN, CH-1211 Geneva 23, Switzerland*

^b*Department of Physics, National University of Singapore, Singapore 119260*

Abstract

We study configurations consisting of a pair of non-extremal black holes in four dimensions, both with the same mass, and with charges of the same magnitude but opposite sign—dipoles, for short. We present such exact solutions for Einstein-Maxwell theory with arbitrary dilaton coupling, and also solutions to the $U(1)^4$ theories that arise from compactified string/M-theory. Despite the fact that the solutions are very complicated, physical properties of these black holes, such as their area, charge, and interaction energy, admit simple expressions. We also succeed in providing a microscopic description of the entropy of these black holes using the ‘effective string’ model, and taking into account the interaction between the effective string and anti-string.

¹roberto.emparan@cern.ch. Also at Departamento de Física Teórica, Universidad del País Vasco, Bilbao, Spain.

1 Introduction

In spite of the enormous recent progress in understanding the microscopic origin of the Bekenstein-Hawking entropy using string theory [1], the criticism is sometimes raised that only a very restricted class of systems are amenable to such a study. Indeed, most of the work in this area has been performed in situations where either supersymmetry is present, or at least the deviation from it is, in some sense, small. In this respect, the exact microscopic calculation of the entropy of the Schwarzschild black hole remains as the most outstanding problem. It would also be desirable to have an understanding of more complicated black hole configurations, and of how the microstates of the black hole react when submitted to external influences.

As a matter of fact, the study of string and brane physics in situations far from a supersymmetric state has become a very active area of research [2]. Non-BPS branes and the closely related brane-antibrane configurations have provided considerable insight into non-perturbative aspects of string/M-theory, such as dualities and tachyon condensation. Nevertheless, it is probably fair to say that the extension to include self-gravity (closed string) effects into the study of such systems lags far behind their open string description. Some steps in this direction have been taken in [3, 4, 5, 6, 7, 8, 9, 10, 11].

In this paper we perform a study of a certain class of black hole configurations (which can be lifted to configurations of black branes), whose features provide a connection to several of the above mentioned issues. Specifically, the systems we study consist of a pair of black holes at a finite distance, with equal masses, and charges of the same magnitude but opposite sign. Since these configurations carry an electric (or magnetic) dipole moment, they are referred to as *diholes*. In Ref. [4], building on the earlier work of [12, 13, 14, 15, 3], a comprehensive study of such configurations was carried out for the case when the black holes are extremal, *i.e.*, the horizons are degenerate and the charge is fixed by the mass. In the present paper, we extend the analysis to the case where the black holes are not extremal, so the horizons are not degenerate, and the charge and mass of each black hole are completely independent parameters. We provide both a macroscopic description, *i.e.*, a study of the solution in classical general relativity, and a string/M-theory microscopic analysis of the entropy of the configuration, that takes into account the interaction between the two black holes. In each of these two veins, we build on previous work and considerably extend it in several directions.

For the macroscopic description, our starting point is a class of solutions to Einstein-Maxwell theory, which was built in [16] using a method devised earlier in [17], and which

describes a pair of black holes with arbitrary mass, charge, and rotation parameters. This very general solution, though, is exceedingly complicated, and is given in terms of algebraically implicit functions. Extracting particular cases from the general solution involves a considerable amount of work, and the analysis of their properties is far from obvious. For this reason, we have restricted ourselves in this paper to the subset of parameters that yield the non-extremal diholes mentioned above. We will be able to find simple expressions for the area, charge, interaction energy, and other physical properties of the configurations. Then, we will extend the solutions to theories with arbitrary dilaton coupling, and also to the theories with four abelian gauge fields (and three massless scalars) that arise in the low energy limit of a variety of compactifications of string/M-theory down to four dimensions.

With these solutions in hand, we will turn to the microscopic description of the dihole configuration. Here we follow Ref. [18], where the entropy of a pair of equally charged, near-extremal black holes, close to the extremal Majumdar-Papapetrou solution, is studied. In our case, we will be further away from a supersymmetric system, since supersymmetry is absent even in the limit where the black holes are extremal. In fact, the gauge and gravitational forces between the black holes never cancel each other, instead they add together. Therefore, even the extremal state reflects in a non-trivial way the existence of an interaction.

Some comments are in order regarding the connection between the non-extremal diholes that are the subject of this paper, and the extremal ones of [12, 14, 4]. The metric for the former is much more complicated than the latter. As could be expected, the extremal dihole is contained in the solutions we study as the limit where the black hole horizons become degenerate. This means that the Bonnor solution [12] is a particular case of the solutions in [16], a point we have explicitly checked. In particular, this should clear any lingering doubt about whether the dihole interpretation of Bonnor's solution, as given in [4] (as opposed to the original interpretation as a singular dipole), is the correct one. As we will see, the extremal limit is a rather subtle one, and this may be one of the reasons why it took so long to correctly identify the nature of these solutions. The same remark applies to the interpretation of the dilatonic dihole solutions constructed in [14] (which are not a black hole-white hole pair, as initially suggested). Finally, it is also worth noting that the new dilatonic solutions we present include, in particular, the non-extremal extension of the Kaluza-Klein monopole-antimonopole solution of [13, 15].

2 Black holes in Weyl coordinates

Axisymmetric solutions, such as the ones we are going to study, are adequately described using Weyl's canonical coordinates. It will be useful to review the description of several known solutions which are particular limiting cases of the black diholes that are the subject of this paper.

For static, electric solutions, a Weyl metric is of the form

$$\begin{aligned} ds^2 &= -f dt^2 + f^{-1} [e^{2\gamma} (d\rho^2 + dz^2) + \rho^2 d\varphi^2], \\ A_\mu &= (A_0, 0, 0, 0). \end{aligned} \quad (1)$$

The functions f , γ and A_0 only depend on ρ and z . We are considering here a solution with an electric potential, but one can construct a magnetic solution by taking A_0 to be the dual magnetic potential.

To start with, let us take the Reissner-Nordström solution, with the familiar metric

$$ds^2 = -\frac{r^2 - 2mr + q^2}{r^2} dt^2 + \frac{r^2 dr^2}{r^2 - 2mr + q^2} + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2), \quad (2)$$

and potential $A_0 = -q/r$. We transform the (r, θ) coordinates into (ρ, z) through

$$\begin{aligned} \rho &= \sqrt{r^2 - 2mr + q^2}, \\ z &= (r - m) \cos \theta. \end{aligned} \quad (3)$$

Then (2) becomes of the form (1) with

$$\begin{aligned} f &= \frac{(R_+ + r_+)^2 - 4(m^2 - q^2)}{(R_+ + r_+ + 2m)^2}, \\ e^{2\gamma} &= \frac{(R_+ + r_+)^2 - 4(m^2 - q^2)}{4R_+ r_+}, \\ A_0 &= -\frac{2q}{R_+ + r_+ + 2m}, \end{aligned} \quad (4)$$

where we have defined

$$\begin{aligned} R_+ &= \sqrt{\rho^2 + \left(z + \sqrt{m^2 - q^2}\right)^2}, \\ r_+ &= \sqrt{\rho^2 + \left(z - \sqrt{m^2 - q^2}\right)^2} \end{aligned} \quad (5)$$

(the notation R_+ , r_+ has been so chosen for consistency with future definitions). Figure 1(a) plots the axis of the solution and the interpretation of the variables R_+ , r_+ . In Weyl's

coordinates, the black holes correspond to ‘rods’ along the axis. For the Schwarzschild black hole the rod’s length is equal to $2m$, but as charge is added the length of the rod shrinks. In the extremal limit $m = q$ the rod contracts to a point, a fact that makes this limit a rather singular one to take in Weyl’s coordinates.




Figure 1: Structure of the axis, and interpretation of $R_{\pm}$, $r_{\pm}$, for the Weyl solutions corresponding to (a) Reissner-Nordström, (b) Israel-Khan neutral two-black hole solution. The black hole horizons correspond to the ‘rods’ marked by the bold lines.

Now let us consider the Israel-Khan solution [19], which describes a set of neutral black holes along the z axis. For the case of two black holes of equal mass m , with the rods centered at $z = \pm k$, the solution is

$$f = \left(\frac{R_+ + r_+ - 2m}{R_+ + r_+ + 2m} \right) \left(\frac{R_- + r_- - 2m}{R_- + r_- + 2m} \right),$$

$$e^{2\gamma} = \left(\frac{\rho^2 + (z + k + m)(z + k - m) + R_+ r_+}{2R_+ r_+} \right) \left(\frac{\rho^2 + (z - k - m)(z - k + m) + R_- r_-}{2R_- r_-} \right) \\ \times \left(\frac{\rho^2 + z^2 - (k - m)^2 + r_+ r_-}{\rho^2 + (z - k - m)(z + k - m) + r_+ R_-} \right) \left(\frac{\rho^2 + z^2 - (k + m)^2 + R_+ R_-}{\rho^2 + (z + k + m)(z - k + m) + R_+ r_-} \right), \quad (6)$$

where now (see Figure 1(b))

$$R_{\pm} = \sqrt{\rho^2 + (z \pm (k + m))^2}, \\ r_{\pm} = \sqrt{\rho^2 + (z \pm (k - m))^2}. \quad (7)$$

In this case we have two rods along the axis, each one of length $2m$, and centers separated by a (coordinate) distance $2k$. The function $\log f$ is indeed the linear superposition of the Newtonian potentials created by two rods of constant mass per unit length. On the other hand, the function γ accounts for the general-relativistic interaction between the black holes. The black holes are of course expected to attract each other, and this is encoded in γ as follows. Suppose that, in a region away from the rods, a non-zero value for

$$\lim_{\rho \rightarrow 0} \gamma = \gamma_0 \quad (8)$$

is obtained as one approaches the axis $\rho = 0$. Then, with the standard choice for the periodicity of φ , *i.e.*, $\varphi \sim \varphi + 2\pi$, there is a conical deficit along the axis equal to

$$\delta = 2\pi(1 - e^{-\gamma_0}). \quad (9)$$

For the particular metric (6), one finds that, outside the rods, $\gamma_0 = 0$, hence there is no conical defect there. Instead, on the segment inbetween the rods the conical angle is

$$\delta = -2\pi \frac{m^2}{k^2 - m^2}, \quad (10)$$

which, being negative, indicates the presence of pressure along the axis (a ‘strut’) that keeps the black holes apart. We are considering the case $k > m$, which corresponds to two non-overlapping rods. If $k = m$ the solution actually describes a single Schwarzschild black hole of mass $2m$, whereas if $k < m$ one obtains again two black holes, but now the roles of k and m are exchanged. Also, the solution for an individual Schwarzschild black hole can be recovered from (6) by performing the shift $z \rightarrow z - k$ and then taking the limit $k \rightarrow \infty$. In this limit, one of the black holes is pushed away to infinity.

Finally, we consider the solution that describes the extremal dihole, *i.e.*, two extremal charged black holes, of the same mass, and charges equal in magnitude but of opposite sign:

$$\begin{aligned} f &= \left[\frac{(R_+ + R_-)^2 - 4m^2 - \frac{k^2}{m^2+k^2}(R_+ - R_-)^2}{(R_+ + R_- + 2m)^2 - \frac{k^2}{m^2+k^2}(R_+ - R_-)^2} \right]^2, \\ e^{2\gamma} &= \left[\frac{(R_+ + R_-)^2 - 4m^2 - \frac{k^2}{m^2+k^2}(R_+ - R_-)^2}{4R_+R_-} \right]^4, \\ A_0 &= -\frac{4mk}{\sqrt{m^2+k^2}} \frac{R_+ - R_-}{(R_+ + R_- + 2m)^2 - \frac{k^2}{m^2+k^2}(R_+ - R_-)^2}, \end{aligned} \quad (11)$$

with, now,

$$R_{\pm} = \sqrt{\rho^2 + (z \pm \sqrt{m^2 + k^2})^2}, \quad (12)$$

see Figure 2. In this case the black holes are extremal, *i.e.*, their horizons are degenerate. Hence each black hole is represented by a point on the axis. For more details about this solution, see [4].




Figure 2: The extremal dihole solution in Weyl coordinates. The extremal black hole horizons correspond now to the black dots.

3 The non-extremal dihole

The class of solutions of Einstein-Maxwell theory that describes two black holes along the axis, with arbitrary mass, charge, and rotation parameters, was presented in [16], following a method devised by Sibgatullin [17]. Such solutions are indeed very complicated, and are given in terms of a number of algebraically implicit functions—a considerable amount of work is required to recover particular cases from them. We are interested in the solution for two non-extremal black holes, of equal mass, and charge of the same magnitude but opposite

sign. Henceforth this solution will be referred to as a non-extremal dihole. In the notation of [16], our choice of parameters is

$$\begin{aligned} m_1 = m_2 = m, \quad q_1 = -q_2 = q, \\ a_1 = a_2 = 0, \quad z_1 = -z_2 = k. \end{aligned} \quad (13)$$

Then, the solution in Weyl's canonical form (1),

$$ds^2 = -f dt^2 + f^{-1} [e^{2\gamma} (d\rho^2 + dz^2) + \rho^2 d\varphi^2], \quad (14)$$

is¹

$$f = \frac{\mathcal{A}^2 - \mathcal{B}^2 + \mathcal{C}^2}{(\mathcal{A} + \mathcal{B})^2}, \quad e^{2\gamma} = \frac{\mathcal{A}^2 - \mathcal{B}^2 + \mathcal{C}^2}{K_0 R_+ R_- r_+ r_-}, \quad A_0 = -\frac{\mathcal{C}}{\mathcal{A} + \mathcal{B}}, \quad (15)$$

where $\mathcal{A}$, $\mathcal{B}$ and $\mathcal{C}$ are given by

$$\begin{aligned} \mathcal{A} &= -(R_+ - R_-)(r_+ - r_-)[(\kappa_+^2 + \kappa_-^2)(m^4 + \kappa_+^2 \kappa_-^2) - 4m^2 \kappa_+^2 \kappa_-^2] \\ &\quad - 2(\kappa_+^2 - \kappa_-^2)(R_+ R_- + r_+ r_-)[m^4 - (\kappa_+ \kappa_-)^2] \\ &\quad + (\kappa_+^2 - \kappa_-^2)(R_+ + R_-)(r_+ + r_-)[m^4 + (\kappa_+ \kappa_-)^2], \\ \mathcal{B} &= 4m(\kappa_+^2 - \kappa_-^2)\kappa_+ \kappa_- [(R_+ + R_- + r_+ + r_-)\kappa_+ \kappa_- - (R_+ + R_- - r_+ - r_-)m^2], \\ \mathcal{C} &= 4kq\kappa_+ \kappa_- [(R_+ - R_- + r_+ - r_-)(\kappa_+^2 - m^2)\kappa_- - (R_+ - R_- - r_+ + r_-)(\kappa_-^2 - m^2)\kappa_+]. \end{aligned} \quad (16)$$

Here, we have defined

$$R_{\pm} \equiv \sqrt{\rho^2 + (z \pm (\kappa_+ + \kappa_-))^2}, \quad r_{\pm} \equiv \sqrt{\rho^2 + (z \pm (\kappa_+ - \kappa_-))^2}. \quad (17)$$

The solution depends on three parameters—in physical terms, the mass, charge and separation between the black holes. We have found it convenient to choose these to be m , κ_+ and κ_- , all of them positive. In terms of the parameters q and k introduced in [16], they are expressed as

$$\kappa_{\pm} = \frac{1}{2} \left(\sqrt{m^2 + k^2 + 2k\sqrt{m^2 - q^2}} \pm \sqrt{m^2 + k^2 - 2k\sqrt{m^2 - q^2}} \right). \quad (18)$$

Conversely,

$$\begin{aligned} k^2 &= \kappa_+^2 + \kappa_-^2 - m^2, \\ kq &= \sqrt{(\kappa_+^2 - m^2)(m^2 - \kappa_-^2)}. \end{aligned} \quad (19)$$

The black hole horizons lie on the axis $\rho = 0$ at $-\kappa_- \leq z \pm \kappa_+ \leq \kappa_-$, see Figure 3.

¹Note that there is a misprint in (3.17) of [16]: $z + m$ should be z . This equation is needed to show that $\omega = 0$, *i.e.*, the $g_{t\varphi}$ component of the metric vanishes.




Figure 3: Structure of the axis, and interpretation of $R_{\pm}$, $r_{\pm}$, (17), and $\kappa_{\pm}$. The black hole horizons correspond to the ‘rods’ marked by the bold lines.

We will restrict the parameters to satisfy $\kappa_- \leq m < \kappa_+$. With this choice one obtains a real solution describing two black holes. Then, as is clear from the figure, κ_+ controls the separation between the black holes, while κ_- is a ‘non-extremality’ parameter. The case $\kappa_+ \leq m < \kappa_-$ actually also describes two black holes, since it can be mapped to the former situation by exchanging $\kappa_+ \leftrightarrow \kappa_-$ and also at the same time $r_+ \leftrightarrow r_-$. These solutions, where κ_+ and κ_- exchange their roles as giving the separation and length of the rods, are therefore redundant and we will not need to consider them separately. One may also ask what happens when the two oppositely charged black holes touch, $\kappa_+ = \kappa_-$. In this case the solution turns out to be a neutral one, since (18) then implies $q = 0$ (hence $\mathcal{C} = 0$), and $\kappa_{\pm} = m$. The solution is in fact a Schwarzschild black hole of mass $2m$. Finally, we remark that neither q nor k are the physical charge or separation between the black holes, but simply parameters that, in the limit of large separation between the holes, approximate these two quantities. The actual physical charge will be determined below, while the proper distance between the black holes does not appear to be too relevant, and at any rate will not be needed in the following.

It still remains to determine K_0 , which is a quantity that can depend on the parameters of the solution but not on the coordinates. Which value one chooses for K_0 is dictated by the structure of conical singularities along the axis. As discussed in the previous section, in a portion of the axis away from the horizons, if the periodicity of φ is $\Delta\varphi$, and $\gamma_0 \equiv \gamma|_{\rho \rightarrow 0}$,

then there will be a conical deficit

$$\delta = 2\pi - \Delta\varphi e^{-\gamma_0}. \quad (20)$$

A direct calculation yields, for the portions inbetween the black holes, and outside them,

$$e^{-\gamma_0} = \frac{K_0^{1/2}}{8\kappa_-^2(\kappa_+^2 - m^2)^2}, \quad |z| < \kappa_+ - \kappa_- \quad (21)$$

$$e^{-\gamma_0} = \frac{K_0^{1/2}}{8\kappa_+^2\kappa_-^2(\kappa_+^2 - \kappa_-^2)}, \quad |z| > \kappa_+ + \kappa_- \quad (22)$$

If we choose to cancel the defect at infinity then we set

$$K_0^{1/2} = 8\kappa_+^2\kappa_-^2(\kappa_+^2 - \kappa_-^2), \quad (23)$$

together with $\Delta\varphi = 2\pi$ (only the product $\Delta\varphi K_0^{1/2}$ is significant). In that case, the conical angle on the axis is negative (*i.e.*, an excess angle, or ‘strut’),

$$\delta_{\text{strut}} = -2\pi \frac{m^2(\kappa_+^2 - m^2) + \kappa_+^2(m^2 - \kappa_-^2)}{(\kappa_+^2 - m^2)^2}. \quad (24)$$

On the other hand, if, instead, we had chosen to remove the singularity from the axis inbetween the black holes, we would have found a conical deficit (a ‘cosmic string’) extending out to infinity,

$$\delta_{\text{string}} = 2\pi \frac{m^2(\kappa_+^2 - m^2) + \kappa_+^2(m^2 - \kappa_-^2)}{\kappa_+^2(\kappa_+^2 - \kappa_-^2)}. \quad (25)$$

In the following we will choose to have a strut singularity inbetween the black holes, hence $\Delta\varphi = 2\pi$, and K_0 as in (23).

The solutions described in the previous section can be seen to be recovered in the following limits:

- Individual non-extremal Reissner-Nordström: this corresponds to infinite separation between the black holes, *i.e.*, $k \rightarrow \infty$, $\kappa_+ \rightarrow k$, with m and $\kappa_- \rightarrow \sqrt{m^2 - q^2}$ finite. In order to leave one of the black holes at a finite distance while the other is sent to infinity, first shift $z \rightarrow z - \kappa_+$, and then take $k \rightarrow \infty$.
- Israel-Khan (two Schwarzschild black holes): $q = 0$, $\kappa_+ = k$, $\kappa_- = m$.
- Extremal dihole: $q = m$, $\kappa_+ = \sqrt{m^2 + k^2}$, $\kappa_- = 0$.

The limit for the extremal dihole is particularly tricky, since the rods along the axis shrink to points, $\kappa_- \rightarrow 0$, and $R_{\pm} \rightarrow r_{\pm}$. Since $\mathcal{A}$, $\mathcal{B}$ and $\mathcal{C}$ all vanish with leading order $O(q - m)^2$, the $q \rightarrow m$ limit must be taken with care. It turns out that the expressions simplify considerably by changing to prolate spheroidal coordinates, (x, y) , with $2x \equiv (R_+ + R_-)/\sqrt{m^2 + k^2}$, $2y \equiv (R_+ - R_-)/\sqrt{m^2 + k^2}$. After lengthy manipulations one then recovers the same expressions as in (11).

There is a last limit of some interest, namely the limit $k = 0$. This implies $\kappa_+ = m$, $\kappa_- = 0$, and the charge vanishes. The solution reduces to the Darms solution, with

$$f = \left(\frac{R_+ + R_- - 2m}{R_+ + R_- + 2m} \right)^2. \quad (26)$$

For this solution, not only do the rods shrink to naked null singularities, but also the segment inbetween them develops an infinite conical defect. It is also the same solution as is obtained in the limit $k = 0$ of the extremal dihole, where its dipole moment vanishes. The significance of this solution in this context is somewhat obscure.

4 Physical properties of the dihole

In spite of its daunting aspect, the physical properties of the solution (15) can be computed in a straightforward manner and will take simple forms in terms of m and $\kappa_{\pm}$. First, note that by examining the solution at asymptotically large distances one quite easily sees that the total mass is $2m$ and the electric dipole moment $2kq$. Other properties, such as the area, temperature and charge of the black holes depend on the form of the solution close to the horizons, and will be calculated now.

Let us start with the computation of the area of the black holes. For definiteness, one can focus on the black hole that lies along the segment $-\kappa_- \leq z - \kappa_+ \leq \kappa_-$, the area of the other one being obviously the same. This area is given by

$$A_{\text{bh}} = \int d\varphi \, dz \sqrt{g_{zz} g_{\varphi\varphi}}|_{\rho=0} = 4\pi\kappa_- \left(\frac{\rho e^{\gamma}}{f} \right)_{\rho=0}. \quad (27)$$

The limit $\rho \rightarrow 0$ has to be taken with a bit of care, but does not present much complication. The calculations are long but straightforward, and one readily finds the pleasantly simple result

$$\left(\frac{\rho e^{\gamma}}{f} \right)_{\rho=0} = \frac{(\kappa_+ + m)^2 (\kappa_- + m)^2}{\kappa_+ \kappa_- (\kappa_+ + \kappa_-)}. \quad (28)$$

Hence the area of each black hole is

$$A_{\text{bh}} = \frac{4\pi (\kappa_+ + m)^2 (\kappa_- + m)^2}{\kappa_+ (\kappa_+ + \kappa_-)} . \quad (29)$$

As a check, the areas for the three limiting cases above (individual Reissner-Nordström black hole, Israel-Khan two Schwarzschild black holes, and extremal dihole) can be readily recovered from here.

The temperature $T = \beta^{-1}$ of the black holes can be calculated, by continuing to Euclidean time, to be

$$\begin{aligned} \beta &= 4\pi \left(\frac{e^\gamma}{\partial_\rho f} \right)_{\rho=0} \\ &= 2\pi \frac{(\kappa_+ + m)^2 (\kappa_- + m)^2}{\kappa_+ \kappa_- (\kappa_+ + \kappa_-)} . \end{aligned} \quad (30)$$

In the above limits it reproduces again the expected results. Note that one obtains the simple relation

$$A_{\text{bh}} = 2\beta\kappa_- , \quad (31)$$

which indeed is a straightforward consequence of (27) and (30).

Next we will compute the charge of each black hole. To do so, we employ Gauss' law $Q = \frac{1}{4\pi} \int_{S^2} *F$, taking the (topological) sphere S^2 to be a surface at constant small ρ closely surrounding the black hole horizon. In that case, it is straightforward to find that the black hole charge is given by

$$\begin{aligned} Q &= \frac{1}{4\pi} \int d\varphi \int_{\kappa_+ - \kappa_-}^{\kappa_+ + \kappa_-} dz \left(\frac{\rho}{f} \partial_\rho A_0 \right)_{\rho=0} \\ &= \frac{\sqrt{(\kappa_+^2 - m^2)(m^2 - \kappa_-^2)}}{\kappa_+ - m} . \end{aligned} \quad (32)$$

Notice that, with (19), this is simply

$$Q = \frac{kq}{\kappa_+ - m} . \quad (33)$$

Later we will also require the value of the electric potential at the horizons, Φ . In order to fix the arbitrary additive constant in the potential A_0 , in (15) we have made the choice that A_0 vanishes at asymptotic infinity. This is a reasonable choice, since in contrast to the situation for an individual black hole, or for a number of equally charged black holes, one cannot

choose the arbitrary constant in the potential to make the potentials vanish simultaneously on both horizons. Then, from (15), it follows that

$$\begin{aligned}\Phi &= \sqrt{\frac{(\kappa_+ - m)(m - \kappa_-)}{(\kappa_+ + m)(m + \kappa_-)}} \\ &= \frac{m - \kappa_-}{Q}.\end{aligned}\tag{34}$$

Q and Φ each change sign from one black hole to the other, but the product $Q\Phi$ has the same sign for both.

Finally, we calculate the interaction energy between the black holes. A straightforward hamiltonian analysis shows that the energy associated to a conical defect δ along the z axis is

$$E_\delta = -\frac{\delta}{4} \int dz \, N \sqrt{g_{zz}}|_{\rho=0},\tag{35}$$

where $N = \sqrt{-g_{tt}}$ is the lapse function for the static metric under consideration. Identifying this energy with the interaction energy between the black holes, $V_{\text{int}} \equiv E_\delta$, we get

$$\begin{aligned}V_{\text{int}} &= \frac{\kappa_+ - \kappa_-}{2} (e^{\gamma_0} - 1) \\ &= -\frac{\kappa_+^2 (\kappa_+^2 - \kappa_-^2) - (\kappa_+^2 - m^2)^2}{2\kappa_+^2 (\kappa_+ + \kappa_-)}.\end{aligned}\tag{36}$$

The value of γ_0 used here is the one obtained from (21) and (23). At large separation, *i.e.*, large κ_+ , the interaction energy becomes

$$V_{\text{int}} \simeq -\frac{m^2 + q^2}{2\kappa_+},\tag{37}$$

which is indeed the result expected for the gravitational and electric interaction of two point masses and charges at a distance $\sim 2\kappa_+$.

The remarkable conclusion of this section is that, even if the metric for the non-extremal dihole is a very complicated one, we have managed to obtain simple expressions for all the physical properties of the configuration.

5 Dilatonic and string/M-theory diholes

5.1 Dilatonic dihole solution

We would now like to generalise the non-extremal dihole solution to a solution of Einstein-Maxwell-dilaton theory:

$$I = \frac{1}{16\pi G} \int d^4x \sqrt{-g} \left(R - 2(\nabla\phi)^2 - e^{-2\alpha\phi} F^2 \right),\tag{38}$$

with arbitrary dilaton coupling α . This is necessary should we want to make contact with Kaluza-Klein theory ($\alpha = \sqrt{3}$) or a particular low-energy effective limit of string theory (the simplest, $\alpha = 1$).

It was shown in [11] how one can generate a static, axisymmetric solution of the Einstein-Maxwell-dilaton equations starting from a stationary, axisymmetric solution of the vacuum Einstein equations. Indeed, this procedure was used in [14] to generate the extreme dilatonic dihole solution (for general α) from the Kerr solution. It is straightforward to modify this procedure to generate a general- α solution starting from the corresponding $\alpha = 0$ (pure Einstein-Maxwell) solution, without any reference to the seed vacuum Einstein solution. We shall now outline this procedure, leaving the reader to fill in the details using the results of [11].

Suppose we have an electrically charged² solution of Einstein-Maxwell theory specified by f , γ and A_0 , as in (1). Then the general dilatonic solution has a new f' , γ' , A'_0 , and dilaton ϕ , given by

$$\begin{aligned} f' &= f^{\frac{1}{1+\alpha^2}} e^{-2\alpha\tilde{\phi}}, \\ \gamma' &= \frac{1}{1+\alpha^2} \gamma + \tilde{\gamma}, \\ A'_0 &= \frac{1}{\sqrt{1+\alpha^2}} A_0, \\ e^{2\phi} &= f^{\frac{\alpha}{1+\alpha^2}} e^{2\tilde{\phi}}, \end{aligned} \tag{39}$$

where $\tilde{\phi}$ is a harmonic function satisfying

$$\frac{\partial^2 \tilde{\phi}}{\partial \rho^2} + \frac{\partial^2 \tilde{\phi}}{\partial z^2} + \frac{1}{\rho} \frac{\partial \tilde{\phi}}{\partial \rho} = 0, \tag{40}$$

and $\tilde{\gamma}$ is given in terms of $\tilde{\phi}$ by

$$\begin{aligned} \frac{\partial \tilde{\gamma}}{\partial \rho} &= (1+\alpha^2) \rho \left[\left(\frac{\partial \tilde{\phi}}{\partial \rho} \right)^2 - \left(\frac{\partial \tilde{\phi}}{\partial z} \right)^2 \right], \\ \frac{\partial \tilde{\gamma}}{\partial z} &= 2(1+\alpha^2) \rho \frac{\partial \tilde{\phi}}{\partial \rho} \frac{\partial \tilde{\phi}}{\partial z}. \end{aligned} \tag{41}$$

This solution-generating procedure can be used, for instance, to obtain the extreme dilatonic dihole solution starting from the Bonnor solution. It turns out that $\tilde{\phi} = 0$ in this case [11]. However, this is not necessarily so in general. For example, we must take

$$e^{-2\tilde{\phi}} = \left(\frac{R_+ + r_+ - 2\sqrt{m^2 - q^2}}{R_+ + r_+ + 2\sqrt{m^2 - q^2}} \right)^{\frac{\alpha}{1+\alpha^2}}, \tag{42}$$

²The same results would apply to a magnetically charged solution, provided we replace A_0 by the magnetic potential A_3 and set $\phi \rightarrow -\phi$. However, this solution-generating procedure fails for dyonic solutions.

if we want to generate the non-extreme dilatonic black hole solution from the Reissner-Nordström solution. In this case, we obtain (in Weyl coordinates)

$$\begin{aligned}
f &= \frac{R_+ + r_+ - 2\sqrt{m^2 - q^2}}{R_+ + r_+ + 2m} \left(\frac{R_+ + r_+ + 2\sqrt{m^2 - q^2}}{R_+ + r_+ + 2m} \right)^{\frac{1-\alpha^2}{1+\alpha^2}}, \\
e^{2\gamma} &= \frac{(R_+ + r_+)^2 - 4(m^2 - q^2)}{4R_+ r_+}, \\
A_0 &= -\frac{1}{\sqrt{1+\alpha^2}} \frac{2q}{R_+ + r_+ + 2m}, \\
e^{2\phi} &= \left(\frac{R_+ + r_+ + 2\sqrt{m^2 - q^2}}{R_+ + r_+ + 2m} \right)^{\frac{2\alpha}{1+\alpha^2}}, \tag{43}
\end{aligned}$$

which indeed corresponds to the dilatonic black hole solution [20].

Applying the solution-generating procedure (39) to the non-extremal dihole solution (15), we obtain

$$\begin{aligned}
ds^2 &= -f dt^2 + f^{-1} [e^{2\gamma} (d\rho^2 + dz^2) + \rho^2 d\varphi^2], \\
A_0 &= -\frac{1}{\sqrt{1+\alpha^2}} \frac{\mathcal{C}}{\mathcal{A} + \mathcal{B}}, \\
e^{2\phi} &= \left[\frac{\mathcal{A}^2 - \mathcal{B}^2 + \mathcal{C}^2}{(\mathcal{A} + \mathcal{B})^2} \right]^{\frac{\alpha}{1+\alpha^2}} e^{2\tilde{\phi}}, \tag{44}
\end{aligned}$$

where

$$f = \left[\frac{\mathcal{A}^2 - \mathcal{B}^2 + \mathcal{C}^2}{(\mathcal{A} + \mathcal{B})^2} \right]^{\frac{1}{1+\alpha^2}} e^{-2\alpha\tilde{\phi}}, \quad e^{2\gamma} = \left[\frac{\mathcal{A}^2 - \mathcal{B}^2 + \mathcal{C}^2}{K_0 R_+ R_- r_+ r_-} \right]^{\frac{1}{1+\alpha^2}} e^{2\tilde{\gamma}}, \tag{45}$$

and $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are the same as in (16). We shall choose the harmonic function $\tilde{\phi}$ to be given by

$$e^{-2\tilde{\phi}} = \left[\left(\frac{R_+ + r_+ - 2\kappa_-}{R_+ + r_+ + 2\kappa_-} \right) \left(\frac{R_- + r_- - 2\kappa_-}{R_- + r_- + 2\kappa_-} \right) \right]^{\frac{\alpha}{1+\alpha^2}}, \tag{46}$$

from which we derive

$$\begin{aligned}
e^{2\tilde{\gamma}} &= \left[\frac{\rho^2 + (z + \kappa_+ + \kappa_-)(z + \kappa_+ - \kappa_-) + R_+ r_+}{2R_+ r_+} \right. \\
&\times \frac{\rho^2 + (z - \kappa_+ - \kappa_-)(z - \kappa_+ + \kappa_-) + R_- r_-}{2R_- r_-} \\
&\times \frac{\rho^2 + z^2 - (\kappa_+ + \kappa_-)^2 + R_+ R_-}{\rho^2 + (z + \kappa_+ + \kappa_-)(z - \kappa_+ + \kappa_-) + R_+ r_-} \\
&\times \left. \frac{\rho^2 + z^2 - (\kappa_+ - \kappa_-)^2 + r_+ r_-}{\rho^2 + (z + \kappa_+ - \kappa_-)(z - \kappa_+ - \kappa_-) + r_+ R_-} \right]^{\frac{\alpha^2}{1+\alpha^2}}, \tag{47}
\end{aligned}$$

using the results of [19]. With this, we have the complete non-extremal dilatonic dihole solution. It correctly reduces to the solution of [14] in the extreme limit. Furthermore, the

individual non-extremal dilatonic black holes (given by (43)) can be recovered in the infinite separation limit, thus confirming that the choice of $\tilde{\phi}$ in (46) is the correct one.

We briefly turn to some physical properties of this solution. Now, we have $e^{\tilde{\gamma}}|_{\rho=0} = 1$ on the axis outside the black holes. To cancel the conical defect at infinity, K_0 therefore has to assume the same value as in the Einstein-Maxwell case, (23). With this value, the area of each black hole is

$$A_{\text{bh}} = \frac{4\pi}{\kappa_+} \frac{(4\kappa_-^2)^{\frac{\alpha^2}{1+\alpha^2}} [(\kappa_+ + m)(\kappa_- + m)]^{\frac{2}{1+\alpha^2}}}{(\kappa_+ + \kappa_-)^{\frac{1-\alpha^2}{1+\alpha^2}}}. \quad (48)$$

Note that when $\alpha \neq 0$, the area vanishes in the extreme limit as expected. It also gives the correct expression in the limit of infinite separation. We can also read off from A_{bh} , the inverse temperature β of each black hole using (31).

The electric charge of each black hole is

$$\begin{aligned} Q &= \frac{1}{4\pi} \int d\varphi \int_{\kappa_+ - \kappa_-}^{\kappa_+ + \kappa_-} dz \left(\frac{\rho}{f} e^{2\alpha\phi} \partial_\rho A_0 \right)_{\rho=0} \\ &= \frac{1}{\sqrt{1+\alpha^2}} \frac{\sqrt{(\kappa_+^2 - m^2)(m^2 - \kappa_-^2)}}{\kappa_+ - m}, \end{aligned} \quad (49)$$

while the value of the potential at the horizon is

$$\Phi = \sqrt{\frac{1}{1+\alpha^2} \frac{(\kappa_+ - m)(m - \kappa_-)}{(\kappa_+ + m)(m + \kappa_-)}}. \quad (50)$$

Lastly, we note that $e^{\tilde{\gamma}}|_{\rho=0} = \left[\frac{\kappa_+^2 - \kappa_-^2}{\kappa_+^2} \right]^{\frac{\alpha^2}{1+\alpha^2}}$ on the axis inbetween the black holes. The interaction energy between the black holes can then be calculated to be

$$V_{\text{int}} = - \frac{\kappa_+^2(\kappa_+^2 - \kappa_-^2) - (\kappa_+^2 - m^2)^{\frac{2}{1+\alpha^2}}(\kappa_+^2 - \kappa_-^2)^{\frac{2\alpha^2}{1+\alpha^2}}}{2\kappa_+^2(\kappa_+ + \kappa_-)}. \quad (51)$$

To find the corresponding magnetically charged dihole solution, we have to perform the duality transformation $F \rightarrow *F$, $\phi \rightarrow -\phi$, of which the former can be written as

$$\frac{\partial A_3}{\partial \rho} = \rho f^{-1} e^{-2\alpha\phi} \frac{\partial A_0}{\partial z}, \quad \frac{\partial A_3}{\partial z} = -\rho f^{-1} e^{-2\alpha\phi} \frac{\partial A_0}{\partial \rho}. \quad (52)$$

Unfortunately, integrating these expressions to obtain the magnetic potential A_3 appears to be a formidable task, and we have not been able to obtain an explicit expression for it. We end off with the remark that the magnetic dihole solution in Kaluza-Klein theory generalises the monopole-antimonopole solution found in [13, 15]. It can also be uplifted to Type IIA string theory, to describe a non-extremal (black) D6-anti-D6-brane configuration, thereby extending the extremal solution in [15, 3].

5.2 $U(1)^4$ dihole solution

In this subsection, we turn to a theory consisting of four abelian gauge fields and three scalar fields, given by the action

$$I = \frac{1}{16\pi G} \int d^4x \sqrt{-g} \left\{ R - \frac{1}{2} [(\partial\eta)^2 + (\partial\sigma)^2 + (\partial\tau)^2] - \frac{e^{-\eta}}{4} [e^{-\sigma-\tau} F_{(1)}^2 + e^{-\sigma+\tau} F_{(2)}^2 + e^{\sigma+\tau} F_{(3)}^2 + e^{\sigma-\tau} F_{(4)}^2] \right\}. \quad (53)$$

This so-called $U(1)^4$ theory is of wide interest because it emerges as a consistent truncation of a number of different compactifications of low-energy string theory and M-theory [21]. The one that we shall focus on below consists of three 5-branes in M-theory intersecting on a common string, along which momentum is flowing [22].

Solutions to this theory describing a pair of non-extremal black holes accelerating apart were constructed in [23]. Now we aim to find the non-extremal dihole solution in this theory. The individual black holes must carry equal and opposite values of four charges under each of the gauge fields, two of which are electric and the other two magnetic. Hence the dihole solution will depend on a total of six parameters: four for each of the charges, one for the total mass, and one for the distance between the black holes. It will be convenient to choose these parameters to be $\kappa_{\pm}$, and m_i , $i = 1, \dots, 4$. Notice that $\kappa_{\pm}$ do not depend on the index i labeling each $U(1)$.

Bearing the latter fact in mind, it is possible to find a simple factorised form for the dihole line element:

$$ds^2 = -(f_1 f_2 f_3 f_4)^{\frac{1}{4}} dt^2 + (f_1 f_2 f_3 f_4)^{-\frac{1}{4}} \left[e^{\frac{1}{2}(\gamma_1 + \gamma_2 + \gamma_3 + \gamma_4)} (d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right], \quad (54)$$

where

$$f_i \equiv \frac{\mathcal{A}_i^2 - \mathcal{B}_i^2 + \mathcal{C}_i^2}{(\mathcal{A}_i + \mathcal{B}_i)^2}, \quad e^{2\gamma_i} \equiv \frac{\mathcal{A}_i^2 - \mathcal{B}_i^2 + \mathcal{C}_i^2}{K_0 R_+ R_- r_+ r_-}. \quad (55)$$

$\mathcal{A}_i$, $\mathcal{B}_i$ and $\mathcal{C}_i$ are defined as in (16), but with m replaced by m_i . Again, K_0 should be chosen as in (23) if one requires regularity on the axis outside the dihole (notice this is consistent with K_0 not having an index i). The three scalar fields are given by

$$e^{-2\eta} = \left(\frac{f_1 f_3}{f_2 f_4} \right)^{\frac{1}{2}}, \quad e^{-2\sigma} = \left(\frac{f_1 f_4}{f_2 f_3} \right)^{\frac{1}{2}}, \quad e^{-2\tau} = \left(\frac{f_1 f_2}{f_3 f_4} \right)^{\frac{1}{2}}. \quad (56)$$

Furthermore, the electric potentials $A_{(2)}$ and $A_{(4)}$ are

$$A_{(i)} = -\frac{\mathcal{C}_i}{\mathcal{A}_i + \mathcal{B}_i}, \quad i = 2, 4, \quad (57)$$

while the magnetic potentials $A_{(1)}$ and $A_{(3)}$ are given implicitly by applying the analogue of the duality transformation (52) on the corresponding electric potentials. We will not need their explicit forms.

This solution has been checked to satisfy the field equations coming from (53). Another consistency check involves setting 1, 2, 3 or 4 of the charges to be non-zero and equal. As is well known, this should reduce to the dilatonic dihole solution (44) for $\alpha = \sqrt{3}$, 1, $\frac{1}{\sqrt{3}}$ or 0, respectively [21]. Indeed, this can readily be checked if we recall that in the chargeless limit $\kappa_- = m_i$,

$$f_i = \left(\frac{R_+ + r_+ - 2\kappa_-}{R_+ + r_+ + 2\kappa_-} \right) \left(\frac{R_- + r_- - 2\kappa_-}{R_- + r_- + 2\kappa_-} \right). \quad (58)$$

This expression is of the right form to reproduce the $e^{-2\alpha\tilde{\phi}}$ terms in the metric for each of the four cases. A similar analysis of the Σ_i terms shows that they correctly give the $e^{2\tilde{\gamma}}$ term in the metric as well.

The total mass of the solution, as measured at infinity, is $\frac{1}{2}\sum_{i=1}^4 m_i$, and the dipole moment for each of the gauge fields is $2kq_i = 2\sqrt{(\kappa_+^2 - m_i^2)(m_i^2 - \kappa_-^2)}$. Finally, the area, charge and potential at the horizon for each of the two black holes, and the interaction energy between them, are

$$A_{\text{bh}} = 4\pi \frac{\prod_{i=1}^4 \sqrt{(\kappa_+ + m_i)(\kappa_- + m_i)}}{\kappa_+(\kappa_+ + \kappa_-)} = 2\beta\kappa_-, \quad (59)$$

$$Q_i = \frac{\sqrt{(\kappa_+^2 - m_i^2)(m_i^2 - \kappa_-^2)}}{\kappa_+ - m_i}, \quad (60)$$

$$\Phi_i = \frac{m_i - \kappa_-}{Q_i}, \quad (61)$$

$$V_{\text{int}} = -\frac{\kappa_+^2(\kappa_+^2 - \kappa_-^2) - \prod_{i=1}^4 \sqrt{\kappa_+^2 - m_i^2}}{2\kappa_+^2(\kappa_+ + \kappa_-)}. \quad (62)$$

6 Statistical mechanics of the dihole

We now want to find a consistent microscopic description, within string/M-theory, of the entropy of the dihole in the case where the two oppositely charged black holes are extremal or nearly extremal.

Our study will be along the lines of the beautiful analysis in [18], where a configuration of two black holes with equal charges of *the same sign* is studied. There is a significant difference between our dihole configuration and the system in [18]. Namely, in the case that the black holes have charges of the same sign, the force between them vanishes in the extremal limit.

Therefore, in that case the near-extremal limit is also one where the interaction between the black holes is small. The ground state of this system is a BPS state. This is not the case for the configuration we study: the force between both black holes does not cancel in the extremal limit, and all the supersymmetries are broken. Thus, we will need to study the extremal dihole first.

We will work in the approximation where the distance between the black holes is large,

$$\kappa_+ \gg m, \kappa_- . \quad (63)$$

In this case, the interaction between the black holes is weak. It is unclear whether the solution will be reliable, for thermodynamic and semiclassical analysis, when the interaction energy that is stored in the strut is of the order of, or larger than, the mass of the black holes. As in [18], we regard the strut as a sort of boundary condition on the system that accounts for the interaction. If the strut singularity becomes too strong, the distortion introduced may invalidate the approach. Recall that, indeed, in the limit $k = 0$ the horizons disappear and are replaced by a naked singularity of diverging curvature.

6.1 Energy and entropy of the extremal dihole

As a preliminary for the microscopic analysis of the near-extremal dihole, we will discuss here the energetics of the configuration when the black holes are extremal, *i.e.*, their horizons are degenerate. For simplicity, we will consider the non-dilatonic case. Recall that this configuration is described by Bonnor's solution, which can be obtained from (15) as the particular case where

$$q = m, \quad \kappa_+ = \sqrt{m^2 + k^2}, \quad \kappa_- = 0 . \quad (64)$$

Hence, the area and charge of each black hole become

$$A_{\text{bh}} = 4\pi m^2 \frac{(m + \sqrt{m^2 + k^2})^2}{m^2 + k^2}, \quad Q = \frac{m}{k}(m + \sqrt{m^2 + k^2}), \quad (65)$$

and the interaction energy between them,

$$V_{\text{int}} = -\frac{m^2}{2} \frac{m^2 + 2k^2}{(m^2 + k^2)^{3/2}} . \quad (66)$$

On the other hand, the *total* energy of the system, E , can be obtained by using, *e.g.*, the formulation in [24], with the result that

$$E = 2m - V_{\text{int}} . \quad (67)$$

Apparently, the sign of the interaction term is the opposite of what might have been expected (recall that $V_{\text{int}} < 0$). This, however, is not really the case. In isolation, the mass of each black hole is equal to its charge. The interaction energy should be equal to the difference between the total energy at finite separation, and the energy when the black holes are infinitely apart, *i.e.*, the BPS mass $2M_{BPS} = 2Q$. In terms of the charge of the black holes, the parameter m is, in the approximation of large separation,

$$m = Q \left(1 - \frac{Q}{k} + O(Q/k)^2 \right). \quad (68)$$

In this approximation we also have

$$V_{\text{int}} = -\frac{m^2}{k} [1 + O(m/k)]. \quad (69)$$

Hence we recover

$$E = 2Q - \frac{Q^2}{k} [1 + O(Q/k)], \quad (70)$$

i.e., to this order, $E = 2M_{BPS} + V_{\text{int}}$, in which the interaction term comes out with the expected sign.

This imposes a consistency check on the interpretation of the area as a measure of black hole internal states. Observe that, by introducing an interaction, the relationships $m(Q)$ and $E(Q)$ get correction terms of order Q/k . Then, in principle one would expect that the entropy $S(Q)$ should also receive corrections at that same order. However, since $E = 2M_{BPS} + V_{\text{int}} + O(Q^3/k^2)$, the internal energy of each black hole remains $E_{\text{bh}} = M_{BPS}[1 + O(Q^2/k^2)]$. That is, the black holes remain in their ground state, to the order Q/k we are working. If the entropy (equal to one quarter of the horizon area) is to be a measure of the number of internal degrees of freedom, then it should remain the same when expressed in terms of the conserved quantity, Q , that measures the number of microstates for the extremal black hole. Hence, the corrections of order Q/k to the entropy must vanish. Indeed, from (65), it follows that this entropy is

$$S = \frac{A_{\text{bh}}}{4} = \pi Q^2 [1 + O(Q/k)^2], \quad (71)$$

as required. It is therefore fully consistent to say that, despite the interaction, and the accompanying distortion of the horizon, the black holes are not excited above their ground state. To see the effect of the interaction on the internal degrees of freedom of the black hole we will have to ‘heat up’ the system above the extremal ground state. This we will do in the next subsection.

It should nonetheless be noted that the relation $A_{\text{bh}} = 4\pi Q^2$ gets spoiled by corrections of order $O(Q/k)^2$. This is perhaps a sign that the solution with the strut ceases to be sensible

as the distance between the black holes decreases³. As mentioned, it becomes singular when $k = 0$. In contrast, if an external axisymmetric field is introduced so as to exactly balance the attraction between the black holes, then the conical singularity disappears, and the relation $A_{\text{bh}} = 4\pi Q^2$ remains an exact one for any value of k [4]. In addition, the solution is non-singular (outside the horizons) even in the limit $k = 0$ —the black holes reach a minimal, non-zero separation, and do not merge. Hence, it appears that by balancing the forces between the black holes with external fields, one obtains a system that is physically sensible for all values of the separation.

6.2 Microscopic entropy of the near-extremal dihole

Just as the individual four-dimensional charged black hole solution, the dihole can be embedded into string and M-theory in a variety of ways. Instead of the embeddings used in [25], it is simpler and more convenient to lift the solution to a configuration of three M5-branes intersecting on a string, with momenta running along the string [22]. If all the directions parallel to the M5-branes, except for their common intersection, are compactified, then one obtains the metric of a five-dimensional string, or in our case, a pair of oppositely charged strings, with momentum running along them in opposite directions—a ‘string-anti-string’ pair. Let us make the choice that $i = 1, 2, 3$ represent the charges associated to each of the M5-branes, and $i = 4$ corresponds to the momentum along the effective string intersection. Explicitly, the metric of the five-dimensional string-anti-string is

$$\begin{aligned} ds^2 = & (f_1 f_2 f_3)^{1/6} \left[-f_4^{1/2} dt^2 + \frac{1}{f_4^{1/2}} (dy - A_{(4)} dt)^2 \right] \\ & + \frac{1}{(f_1 f_2 f_3)^{1/3}} \left[e^{(\gamma_1 + \gamma_2 + \gamma_3 + \gamma_4)/2} (d\rho^2 + dz^2) + \rho^2 d\varphi^2 \right], \end{aligned} \quad (72)$$

where $A_{(4)}$, f_i , γ_i are those in (55) and (57).

It may also be useful to give the solution for the extremal limit. In this case it is more convenient to use the coordinates and notation of [6], in terms of which

$$\begin{aligned} ds^2 = & (T_1 T_2 T_3)^{1/3} \left[-T_4 dt^2 + \frac{1}{T_4} \left(dy + \frac{2k_4 m_4 \cos \theta}{\Sigma_4} dt \right)^2 \right. \\ & \left. + T_4 \frac{\Sigma_1 \Sigma_2 \Sigma_3 \Sigma_4}{(r^2 - \kappa_+^2 \cos^2 \theta)^3} \left(\frac{dr^2}{\Delta} + d\theta^2 \right) \right] + \frac{\Delta \sin^2 \theta}{(T_1 T_2 T_3)^{2/3}} d\varphi^2, \end{aligned} \quad (73)$$

where now $\Delta = r^2 - \kappa_+^2$, $\Sigma_i = (r + m_i)^2 - k_i^2 \cos^2 \theta$, $T_i = (\Delta + k_i^2 \sin^2 \theta)/\Sigma_i$. In the extremal limit, the f_i become $f_i = T_i^2$.

³Notice, though, that the relation $A_{\text{bh}} = 4\pi Q^2$ is neither an exact one for an extremal black hole pierced by a cosmic string. This point deserves further study.

We will now analyze the semiclassical thermodynamics of the four-dimensional $U(1)^4$ dihole, and then, use the effective string model to find a microscopic explanation for the entropy. Again, we are interested in the situation for large distance between the holes, *i.e.*,

$$\kappa_-, m_i \ll \kappa_+. \quad (74)$$

We will also take the ‘dilute gas’ limit for the effective string description of the black holes near extremality, *i.e.*,

$$\kappa_-, m_4 \ll m_1, m_2, m_3. \quad (75)$$

Recall that κ_- plays the role of a non-extremality parameter, the extremal limit being one where $\kappa_- = 0$, while the m_i remain finite.

In order to determine the thermodynamic properties of the dihole, we compute the action for the four-dimensional solution continued to the Euclidean section, $t \rightarrow i\tau$. The Euclidean action can be completely reduced to a calculation of surface terms. The calculations are standard and straightforward. The geometric parts of it yield $\frac{\beta}{2} \sum_i m_i - \frac{1}{4} A_{\text{tot}}$, (here A_{tot} is the sum of the areas of the horizons), plus a contribution from the strut singularity, $-\beta V_{\text{int}}$. The remaining contribution comes from the gauge fields. For the dihole, the electric potentials cannot be made to vanish simultaneously on both horizons. One can instead choose the potential to vanish at infinity, and compute the contributions to the action arising from the (gauge) singularities of the potential A_0 at the horizons. In that case, each horizon gives a contribution $-\frac{\beta}{4} \sum_i Q_i \Phi_i$. Even if the charges have opposite signs for each of the two black holes, the product $Q_i \Phi_i$ has the same sign for both. We are interested in fixing the potentials, and not the charges, at the horizons. For the magnetic fields, this requires the addition of extra boundary terms to the action. Then, Φ_i is the dual magnetic potential. With the appropriate boundary terms, the situations with electric or magnetic charges are dual to each other, as described in [26]. Adding up all the contributions one obtains

$$I = \beta \left(\frac{1}{2} \sum_{i=1}^4 m_i - \frac{1}{2} \sum_{i=1}^4 Q_i \Phi_i - V_{\text{int}} \right) - \frac{1}{4} A_{\text{tot}}. \quad (76)$$

In the long distance approximation,

$$V_{\text{int}} = -\frac{1}{4\kappa_+} \sum_{i=1}^3 Q_i^2 - \frac{m_4^2 - 2\kappa_-^2}{4\kappa_+} + O(m^3/\kappa_+^2). \quad (77)$$

Also, since $m_i - Q_i \Phi_i = \kappa_-$, and $A_{\text{tot}} = 4\beta\kappa_-$, we finally find

$$\begin{aligned} I &= \beta(\kappa_- - V_{\text{int}}) \\ &= \beta \left[\kappa_- + \frac{1}{4\kappa_+} \left(-2\kappa_-^2 + \sum_{i=1}^4 m_i^2 \right) + O(m^3/\kappa_+^2) \right]. \end{aligned} \quad (78)$$

This action provides the grand-canonical potential $W[T, \Phi_i] = I/\beta = E - TS_{\text{tot}} - Q\Phi$. Here S_{tot} is the sum of the entropy of both black holes, which we identify from their area, and $Q\Phi = \sum_{i=1}^4 \frac{1}{2} Q_i \Phi_i$.⁴ The total energy follows as

$$E = \frac{1}{2} \sum_{i=1}^4 m_i + \frac{1}{4\kappa_+} \left(-2\kappa_-^2 + \sum_{i=1}^4 m_i^2 \right) + O(m^3/\kappa_+^2). \quad (79)$$

In the limit (75), and for large κ_+ , we approximate

$$m_i = Q_i \left(1 - \frac{Q_i}{\kappa_+} + O(Q/\kappa_+)^2 \right), \quad i = 1, \dots, 3, \quad (80)$$

so,

$$E = \frac{1}{2} \sum_{i=1}^3 Q_i - \frac{1}{4\kappa_+} \sum_{i=1}^3 Q_i^2 + \frac{m_4}{2} + \frac{m_4^2 - 2\kappa_-^2}{4\kappa_+} + O(m^3/\kappa_+^2). \quad (81)$$

Here we see that, once we have expressed the energy in terms of the M5-brane charges $Q_{1,2,3}$, the interaction energy between the M5-branes (the second term in the right hand side of (81)) comes out with the expected negative sign. This is just as in the extremal case studied above. But now we still have to account for the pieces for the energy and interactions between the excitations along the strings, *i.e.*, the third and fourth terms in the right hand side of (81). To this effect, we have to distinguish carefully between the energy of excitation of each string above the BPS state, call it δE , and the interaction energy between the string-anti-string pair, V_{int} . The BPS ground state of the system consists of the two infinitely separated M5-brane intersections, with the M5-branes in their ground state. The energy of each intersecting brane configuration is $M_{BPS} = \frac{1}{4}(Q_1 + Q_2 + Q_3)$. The total energy must be the sum of three contributions,

$$E = 2(M_{BPS} + \delta E) + V_{\text{int}}. \quad (82)$$

From (81), (82), and (77), we can now identify

$$\delta E = \frac{m_4}{4} \left(1 + \frac{m_4}{\kappa_+} \right) - \frac{\kappa_-^2}{2\kappa_+} + O(m^3/\kappa_+^2). \quad (83)$$

This is the total energy carried by the lightlike left- and right-movers along each of the two antiparallel effective strings,

$$\delta E = P_L + P_R. \quad (84)$$

⁴Properly, one should derive this from W using standard thermodynamics. It can be checked that, within the approximations we take, the entropy-area relation holds. Also, one usually fixes the electric potential at infinity, not on the horizons. It is perhaps possible to, more rigorously, define another thermodynamic ensemble that fixes, *e.g.*, the electric field (which is conjugate to the electric dipole moment). Nevertheless, the present approach leads to the correct results.

To obtain the separate values of P_L and P_R we need the total momentum along each string, $P = P_L - P_R$. This is given by the charge associated to translations along the string direction,

$$P = \frac{Q_4}{4} = \frac{\sqrt{m_4^2 - \kappa_-^2}}{4} \left(1 + \frac{m_4}{\kappa_+} + O(m^2/\kappa_+^2) \right). \quad (85)$$

At this point, it is perhaps convenient to introduce the usual boost parameter δ , such that⁵

$$\begin{aligned} m_4 &= \kappa_- \cosh 2\delta, & m_4 + \kappa_- &= 2\kappa_- \cosh^2 \delta, \\ \sqrt{m_4^2 - \kappa_-^2} &= \kappa_- \sinh 2\delta, & m_4 - \kappa_- &= 2\kappa_- \sinh^2 \delta. \end{aligned} \quad (86)$$

Then, the momentum carried by left and right movers, $P_{L,R} = (\delta E \pm P)/2$, can be expressed as

$$\begin{aligned} P_L &= \frac{\kappa_-}{8} e^{2\delta} \left(1 + \frac{m_4}{\kappa_+} - 2 \frac{\kappa_-}{\kappa_+} e^{-2\delta} + O(m^2/\kappa_+^2) \right), \\ P_R &= \frac{\kappa_-}{8} e^{-2\delta} \left(1 + \frac{m_4}{\kappa_+} - 2 \frac{\kappa_-}{\kappa_+} e^{2\delta} + O(m^2/\kappa_+^2) \right). \end{aligned} \quad (87)$$

With these, we can express the entropy of the dihole in terms of the M5-brane charges and the momentum of the excitations of the string. From (59), we obtain, for each black hole,

$$S = \pi \sqrt{m_1 m_2 m_3} \sqrt{m_4 + \kappa_-} \left(1 - \frac{\kappa_-}{\kappa_+} + \frac{1}{2} \frac{\sum_{i=1}^4 m_i}{\kappa_+} + O(m^2/\kappa_+^2) \right), \quad (88)$$

the total entropy being twice this value. Putting together (80), (87), and (88), we find that, up to the order of approximation we are considering,

$$S = \frac{2\pi}{\sqrt{G}} \sqrt{Q_1 Q_2 Q_3} \left(\sqrt{P_L} + \sqrt{P_R} \right). \quad (89)$$

We have reinstated here the four-dimensional Newton's constant G , which until now had been set to one. The charges $Q_{1,2,3}$ are directly proportional to the numbers $N_{1,2,3}$ of M5-branes. The precise relationship is such that

$$Q_1 Q_2 Q_3 = \frac{GL}{2\pi} N_1 N_2 N_3, \quad (90)$$

where L is the length along the effective string (the length of the compact fifth dimension at asymptotic infinity). On the other hand, the momentum along the string is quantized in the usual way,

$$P_{L,R} = \frac{2\pi}{L} N_{L,R}. \quad (91)$$

⁵Note, however, that the dihole solution with $P \neq 0$ cannot be obtained by boosting the $P = 0$ solution along the direction of the string.

Thus, the Bekenstein-Hawking entropy that we have calculated for each black hole is

$$S = 2\pi\sqrt{N_1N_2N_3}\left(\sqrt{N_L} + \sqrt{N_R}\right). \quad (92)$$

This is precisely the same as that obtained from the statistical mechanics of a $1 + 1$ dimensional field theory with central charge $c = 6N_1N_2N_3$ and $N_{L,R}$ left- and right-moving excitations.

Hence, the effective string model can successfully reproduce, in a microscopic calculation, the entropy of the interacting system of a near-extremal dihole. In string theory terms, the interaction between the two antiparallel effective strings is due to the exchange of closed strings between them. At large separations, this exchange is dominated by the massless states of the closed string. These allow for a description in terms of a classical, Coulombian and Newtonian, static interaction potential, and the effect has been accounted for in a simple way by a shift in the energy levels of each effective string, (82). At shorter distances (of order the string length) the massive modes of the string should become relevant, and eventually a tachyonic instability must appear. This is beyond the approximations herewith considered. We shall simply note that, for the extremal case, a description of the condensation of this tachyon within the supergravity picture has been proposed in [8]. It might be interesting to generalize this beyond the extremal limit.

7 Conclusions and outlook

The excitation of the black holes in the dihole above extremality has resulted in an unexpectedly (to us) complicated solution. This is in contrast to the individual black hole solutions, or the C-metric solutions for pairs of black holes that accelerate apart, where the non-extremal solutions are only slightly more involved than their extremal counterparts. It is somewhat surprising that the extremal solution is so much simpler, since in this case the extremal state does not have any of the special properties, such as supersymmetry or cancellation of forces, that are at work in other configurations. Fortunately, near the horizons the structures simplify sufficiently so that the physical properties of the black holes take simple, manageable forms.

This simplification near the horizons is also related to the reason why the microscopic analysis in terms of an effective string does work. At large separation, the distortion of the extremal horizons caused by the interaction between the black holes is sufficiently small to identify the state as a form of ‘near-BPS’ limit. Above this state, we have added the thermal

excitations of a dilute gas of left and right movers, and the effective string picture appears to be still reliable. We remain in a controlled situation slightly above the supersymmetric state, but one that goes beyond the individual near-extremal black holes or the near-Majumdar-Papapetrou solutions of [18].

By going beyond the extremal limit, and considering the non-extremal diholes, we have been able to exhibit features of the microstates of the black holes that would not have appeared otherwise. Nevertheless, in the microscopic analysis we have needed to make a restriction to large separation between the black holes, so as to obtain a weak strut singularity. It is likely that a better behaved situation is obtained by balancing the dihole (*i.e.*, cancelling the conical singularities) by adding an external field, as was done in [4] for the extremal dihole. It was found there that when the dihole is exactly balanced, the distortion of the horizons completely disappears, in the sense that they become exact, spherically symmetric Bertotti-Robinson throats⁶. Clearly, it should also be possible to similarly balance a non-extremal dihole. Perhaps in this way one can obtain a description which is not subject to the restriction of large separation, eqn. (63). Notice also that this possibility of balancing the forces does not appear to be available to the equal-charge non-extremal configurations of [18]. Unfortunately, it appears that in order to construct the dihole solution in a background field, the magnetic solutions need to be considered first, and the explicit form of the potential A_φ for the magnetic dihole is required. This we have not been able to find for the non-extremal dihole.

Besides the interest in extending the microscopic calculations of the entropy to more complicated systems, or in obtaining new supergravity solutions for brane-anti-brane configurations, there is another, perhaps more speculative motivation for studying black hole/anti-black hole systems. In [27], a mysterious formula for the entropy of the generic $U(1)^4$ black hole, which includes also the neutral Schwarzschild case, was found, which suggested an interpretation in terms of brane-antibrane pairs. While the significance of such a formula remains unclear, one might expect to throw some light on it by studying black hole/anti-black hole configurations. In this respect, notice that the coincidence limit for the non-extremal dihole is better behaved than for the extremal one. This line remains to be pursued.

Another direction in which this work can be extended is by adding rotation to the black holes, either parallel or antiparallel. Since the microscopic degrees of freedom responsible for the rotation add qualitatively new dynamics [28], it would be interesting to see how they are

⁶This is for the non-dilatonic case. In the dilatonic cases the horizons are actually null singularities, but, when balanced, the geometries near these singularities also become spherically symmetric.

affected by the interaction. One can also envisage adding a net charge to the configuration [11]. Work on some of these directions is currently being pursued, and will be reported elsewhere.

Noted added in proof

Fay Dowker (informed, in turn, by David Kastor) has pointed out to us that the extremal dihole in Einstein–Maxwell theory was previously found by Chandrasekhar and Xanthopoulos [29], albeit in a different set of coordinates. While they correctly identified the solution as describing a pair of oppositely charged extremal Reissner–Nordström black holes joined by a string, they did not realise the equivalence of their solution to the Bonnor dipole. They also discussed an apparent violation of the extremality bound, but our analysis in Sec. 6.1 settles this point.

It has also come to our attention that Galtsov et al. [30] also discovered the dilatonic generalisation of the Bonnor solution, independently of Davidson and Gedalin [14]. However, the dihole interpretation of the Bonnor dipole solution apparently remained unknown to them, as with other authors, until the work of [4].

Acknowledgments

ET thanks Yeong Cherng Liang for useful discussions and also for his assistance, in particular, in detecting the misprint highlighted in Footnote 1. RE acknowledges partial support from UPV grant 063.310-EB187/98 and CICYT AEN99-0315.

References

- [1] A. Strominger and C. Vafa, Phys. Lett. B **379** (1996) 99 [hep-th/9601029]. For a review and references, see A. W. Peet, “TASI lectures on black holes in string theory,” hep-th/0008241.
- [2] A. Sen, “Non-BPS states and branes in string theory,” hep-th/9904207.
- [3] A. Sen, JHEP **9710** (1997) 002 [hep-th/9708002].
- [4] R. Emparan, Phys. Rev. D **61** (2000) 104009 [hep-th/9906160].
- [5] D. Youm, Nucl. Phys. B **573** (2000) 223 [hep-th/9908182].

- [6] A. Chattaraputi, R. Emparan and A. Taormina, Nucl. Phys. B **573** (2000) 291 [hep-th/9911007].
- [7] Y. Lozano, Phys. Lett. B **487** (2000) 180 [hep-th/0003226].
- [8] P. Brax, G. Mandal and Y. Oz, Phys. Rev. D **63** (2001) 064008 [hep-th/0005242].
- [9] M. Bertolini, P. Di Vecchia, M. Frau, A. Lerda, R. Marotta and R. Russo, Nucl. Phys. B **590** (2000) 471 [hep-th/0007097].
- [10] P. Bain, JHEP **0104** (2001) 014 [hep-th/0012211].
- [11] Y. C. Liang and E. Teo, Phys. Rev. D **64** (2001) 024019 [hep-th/0101221].
- [12] W. B. Bonnor, Z. Phys. **190** (1966) 444.
- [13] D. J. Gross and M. J. Perry, Nucl. Phys. B **226** (1983) 29.
- [14] A. Davidson and E. Gedalin, Phys. Lett. B **339** (1994) 304 [gr-qc/9408006].
- [15] F. Dowker, J. P. Gauntlett, G. W. Gibbons and G. T. Horowitz, Phys. Rev. D **53** (1996) 7115 [hep-th/9512154].
- [16] V. S. Manko, J. Martín and E. Ruiz, J. Math. Phys. **35** (1994) 6644.
- [17] N. R. Sibgatullin, *Oscillations and Waves in Strong Gravitational and Electromagnetic Fields* (Nauka, Moscow, 1984; English Translation: Springer-Verlag, Berlin, 1991).
- [18] M. S. Costa and M. J. Perry, Nucl. Phys. B **591** (2000) 469 [hep-th/0008106].
- [19] W. Israel and K. A. Khan, Nuovo Cim. **33** (1964) 331.
- [20] G. W. Gibbons and K. Maeda, Nucl. Phys. B **298** (1988) 741. D. Garfinkle, G. T. Horowitz and A. Strominger, Phys. Rev. D **43** (1991) 3140 [Erratum-ibid. D **45** (1992) 3888].
- [21] M. Cvetič and D. Youm, Phys. Rev. D **53** (1996) 584 [hep-th/9507090]. M. Cvetič and A. A. Tseytlin, Phys. Rev. D **53** (1996) 5619 [Erratum-ibid. D **55** (1997) 3907] [hep-th/9512031]. J. Rahmfeld, Phys. Lett. B **372** (1996) 198 [hep-th/9512089]. M. Cvetič and D. Youm, Nucl. Phys. B **472** (1996) 249 [hep-th/9512127].

- [22] I. R. Klebanov and A. A. Tseytlin, Nucl. Phys. B **475** (1996) 179 [hep-th/9604166].
M. Cvetič and A. A. Tseytlin, Nucl. Phys. B **478** (1996) 181 [hep-th/9606033].
- [23] R. Emparan, Nucl. Phys. B **490** (1997) 365 [hep-th/9610170].
- [24] S. W. Hawking and G. T. Horowitz, Class. Quant. Grav. **13** (1996) 1487 [gr-qc/9501014].
- [25] J. M. Maldacena and A. Strominger, Phys. Rev. Lett. **77** (1996) 428 [hep-th/9603060].
C. V. Johnson, R. R. Khuri and R. C. Myers, Phys. Lett. B **378** (1996) 78 [hep-th/9603061].
- [26] S. W. Hawking and S. F. Ross, Phys. Rev. D **52** (1995) 5865 [hep-th/9504019].
- [27] G. T. Horowitz, J. M. Maldacena and A. Strominger, Phys. Lett. B **383** (1996) 151 [hep-th/9603109].
G. T. Horowitz, D. A. Lowe and J. M. Maldacena, Phys. Rev. Lett. **77** (1996) 430 [hep-th/9603195].
- [28] J. C. Breckenridge, D. A. Lowe, R. C. Myers, A. W. Peet, A. Strominger and C. Vafa, Phys. Lett. B **381** (1996) 423 [hep-th/9603078].
- [29] S. Chandrasekhar and B. C. Xanthopoulos, Proc. Roy. Soc. Lond. A **423** (1989) 387.
- [30] D. V. Galtsov, A. A. Garcia and O. V. Kechkin, Class. Quant. Grav. **12** (1995) 2887 [hep-th/9504155].